

Instabilities of bulk fields and anomalies on orbifolds

S. Groot Nibbelink ^{a,1}, H.P. Nilles ^{a,2}, M. Olechowski ^{a,b,3}

^a *Physikalisches Institut der Universität Bonn,
Nussallee 12, 53115 Bonn, Germany.*

^b *Institute of Theoretical Physics, Warsaw University,
Hoża 69, 00-681 Warsaw, Poland.*

Abstract

Bulk matter modes of higher dimensional models generically become unstable in the presence of additional matter multiplets at the branes. This quantum instability is driven by localized Fayet–Iliopoulos terms that attract the bulk zero modes towards the boundary branes. We study this mechanism in the framework of a 5 dimensional $S^1/\mathbb{Z}_2$ orbifold and give conditions for the various possibilities of localization of (chiral) zero modes. This mechanism is quite relevant for realistic model building, as the standard model contains $U(1)$ hypercharge with potentially localized FI–terms. The analysis is closely related to localized anomalies in higher dimensional gauge theories. Five dimensional gauge invariance of the effective action determines the anomaly constraints and fixes the normalization of Chern–Simons terms. The localization of the bulk modes does not effect the anomaly cancellation globally, but the additional heavy Kaluza–Klein modes of the bulk fields may cancel the Chern–Simons terms. We discuss also the potential appearance of the parity anomaly that might render the construction of some orbifold models inconsistent.

¹ E-mail: nibblink@th.physik.uni-bonn.de

² E-mail: nilles@th.physik.uni-bonn.de

³ E-mail: olech@th.physik.uni-bonn.de

1 Introduction

There has recently been a large interest in field theory models of extra dimensions with boundaries. This opens up the possibility that within a given scheme one might have fields living in various dimensions, which may interact with each other in complicated though fascinating ways. The ultimate picture is that such models are presumably part of a low-energy description of string or M-theory, and for that reason it is not surprising that some amount of supersymmetry is present in these extra dimensions. As supersymmetry in higher dimensions corresponds to extended supersymmetry in an effective 4 dimensional theory, some if not all of those supersymmetries should be broken to give us a phenomenologically interesting model that may incorporate the Standard Model of particle physics. Boundaries of, or branes in, extra dimensions lead naturally to a certain amount of supersymmetry breaking and often also to a chiral spectrum in the effective 4 dimensional description.

In the present paper we would like to study two important aspects of theories in extra dimensions, namely their consistency and stability. We consider 5 dimensional supersymmetric field theories, with the extra dimension compactified on the orbifold $S^1/\mathbb{Z}_2$. The orbifold condition naturally leads to boundaries that break half of the supersymmetries. At those branes additional matter may be situated that form well-defined representations of the remaining supersymmetries. For simplicity we consider only the possibility that chiral multiplets (containing chiral fermions and complex scalars) are present on the branes. In the 5 dimensional bulk we assume that a set of hyper multiplets (each consisting of two complex scalars and a Dirac fermion), and vector multiplets (each with 5 dimensional gauge field, a real scalar and a Dirac fermion) are situated.

There can be potential instabilities in models with an arbitrary distribution of matter fields over the bulk and the branes, which indicate that some setups are extremely disfavored. In particular for a $U(1)$ super Yang-Mills-theory in the bulk, we observed in [1] that the divergent Fayet-Iliopoulos (FI)-terms [2] localized at the branes [3, 4] can lead to localization of the zero modes of bulk hyper multiplets. (Localization has been considered in the past, see for example [5]. Consequences of Fayet-Iliopoulos terms on the shape of zero modes have also been studied at the purely classical level in [6, 7].) This dynamical mechanism tells us that certain configurations are unstable, and may therefore not be the appropriate starting point for a thorough phenomenological investigation of such models. In this paper we take a closer look at this phenomenon of localization: in particular we would like to understand which forms of the localized modes are possible and under which conditions they occur.

An even more serious effect signaling instabilities, if not worse, of a quantum theory are anomalies. In [8] it was shown that anomalies in 5 dimensional theories can occur due to bulk fields, but they are situated at the 4 dimensional boundaries only. (Using similar methods anomalies on the orbifold $S^1/\mathbb{Z}_2 \times \mathbb{Z}'_2$ have been discussed in [4, 9, 10].) One question that naturally comes up, is whether models with bulk and brane matter can always be made consistent by an appropriate Chern-Simons term. Even if this is possible, it seems that the consistency of the low-energy spectrum depends on such high-energy interactions: this goes somewhat against the intuition that for anomalies only the low-energy spectrum (the zero modes) are relevant. In addition, does the localization of bulk fermions have consequences for the description of anomalies? In this work we address these questions and show that they are related in an interesting way.

Another serious issue we raise in this paper concerns the very definition of orbifold models with fermions: the orbifold $S^1/\mathbb{Z}_2$ is defined by dividing out a parity in the 5th dimension. What happens if this parity is anomalous? We investigate whether a counter term can render the definition of a model to be consistent.

To achieve these goals we have organized the paper as follows: In section 2 we review properties of five dimensional globally supersymmetric theories on $S^1/\mathbb{Z}_2$. We consider the following bulk multiplets: a $U(1)$ vector multiplet V and a set of charged hyper multiplets H with charge operator Q . In addition we allow for a set of chiral multiplets C_0, C_π with charge matrices q_0, q_π on the branes at $y = 0$ and $y = \pi R$, respectively. Following the arguments of ref. [11] the induced brane supersymmetry transformations of bulk fields are discussed. The consequences of this reasoning for possible FI-terms on branes are confirmed in section 3 by an explicit calculation of the relevant tadpoles that may appear at one-loop. The resulting effective potential is investigated in section 4. In particular we investigate the conditions under which supersymmetric and $U(1)$ gauge symmetry preserving background field configurations do exist.

Next, we describe conditions for localization effects of zero modes to occur in section 5. A careful analysis is presented to classify the possible shapes of the zero modes. This information is accumulated in table 2. A discussion on the mass spectrum of the non-zero modes concludes the section on localization.

The final topic of this paper are anomalies. We first point out that the parity anomaly on the circle S^1 may complicate the definition of the fermionic field theory on the orbifold $S^1/\mathbb{Z}_2$. After that, an alternative derivation of bulk field anomalies is given using topological methods [12]. For consistency the local five dimensional gauge invariance of the orbifold model is essential, we work out the consequences for the matter content and interaction structure of such models. We show that the localization of chiral zero modes changes the description of the local anomalies, but do not affect the consistency of the models.

2 Bulk and (induced) brane supersymmetry

In the bulk we assume that one vector multiplet V and a set of hyper multiplets $H = \{H_b, b = 1, \dots, n\}$ exist. We use conventions and normalizations from [13] for these multiplets, except the auxiliary fields for which we included a factor $-\frac{1}{2}$ to comply with the convention often used in the literature [11]. A $U(1)$ vector multiplet $V = (A_M, \chi, \Phi, \vec{D})$ in five dimensions transforms under supersymmetry as

$$\begin{aligned}\delta A_M &= i\bar{\varepsilon}\gamma_M\chi, & \delta\Phi &= i\bar{\varepsilon}\chi, & \delta\vec{D} &= \bar{\varepsilon}\vec{\sigma}\gamma^M\partial_M\chi, \\ \delta\chi &= \frac{1}{4}F^{MN}\gamma_{MN}\varepsilon - \frac{1}{2}\gamma^M\partial_M\Phi\varepsilon - \frac{1}{2}i\vec{\sigma}\cdot\vec{D}\varepsilon.\end{aligned}\tag{1}$$

Here A_M is a five dimensional gauge field, F_{MN} is the corresponding field strength. The gaugino χ and the supersymmetry parameter ε are symplectic Majorana. As long as the $N = 1$ supersymmetry in five dimensions is manifest, it is convenient to use symplectic Majorana and related reality conditions. In appendix A we collected our conventions of notation and give these reality conditions. To complete the description of the content of the vector multiplet, there is an iso-triplet of auxiliary scalars $\vec{D}$. The Lagrangian for the vector multiplet reads

$$L_V = -\frac{1}{4}F_{MN}F^{MN} + \frac{1}{2}(\partial_M\Phi)^2 + \frac{1}{2}\vec{D}^2 + i\bar{\chi}\gamma^M D_M\chi.\tag{2}$$

A hyper multiplet consists of a set of 4 real scalars h (which are represented as a quaternion), a symplectic Majorana spinor ζ (but with slightly different reality condition than the gauginos, see

appendix A), and a quaternion of 4 real auxiliary fields F . The supersymmetry transformations of the charge hyper multiplets $H = (h, \zeta, F)$ take the form

$$\begin{aligned}\delta h &= i\bar{\varepsilon}\zeta, & \delta F &= i\bar{\varepsilon}\gamma^M D_M \zeta + 2g_5 \bar{\varepsilon} Q h \chi + g_5 \bar{\varepsilon} \Phi Q \zeta, \\ \delta \zeta &= -\gamma^M D_M h \varepsilon - F \varepsilon - i g_5 \Phi Q h \varepsilon, & D_M &= \partial_M + i g_5 A_M Q,\end{aligned}\tag{3}$$

with g_5 the five dimensional gauge coupling and Q the $U(1)$ charge operator. The Lagrangian for the hyper multiplets is given by

$$\begin{aligned}L_H &= \text{tr} \left[\frac{1}{2} D_M h^\dagger D^M h + \frac{1}{2} F^\dagger F - \frac{1}{2} g_5^2 \Phi^2 h^\dagger Q^2 h - \frac{1}{2} g_5 h^\dagger Q h \vec{\sigma} \cdot \vec{D} \right] \\ &\quad + \frac{i}{2} \bar{\zeta} \gamma^M D_M \zeta + \frac{1}{2} g_5 \Phi \bar{\zeta} Q \zeta + 2g_5 \bar{\zeta} Q h \chi.\end{aligned}\tag{4}$$

The trace here is over the $USp(2)$ indices. As the bulk is the interval $S^1/\mathbb{Z}_2$ times four dimensional Minkowski space, orbifold boundary conditions have to be given.

Because of the orbifolding conditions for the fermions, it is not possible to preserve the full five dimensional supersymmetry structure (the 8 super charges). However, the orbifolding preserves 4 global super charges [14]: $\varepsilon = \varepsilon(y) = \varepsilon(-y) = \sigma_3 i \gamma^5 \varepsilon(y) = \sigma_3 i \gamma^5 \varepsilon$. By requiring, in addition, that the gauge symmetry is unbroken, the orbifolding conditions are uniquely defined up to similarity transformations by stating them for the fermions. Concretely, we have for the boundary conditions

$$\begin{aligned}\chi(-y) &= \sigma_3 i \gamma^5 \chi(y), & h(-y) &= -(\mathbb{1} \otimes \sigma_3) h(y) \sigma_3, & \zeta(-y) &= (\mathbb{1} \otimes \sigma_3) i \gamma^5 \zeta(y), \\ \vec{\sigma} \cdot \vec{D}(-y) &= \sigma_3 \vec{\sigma} \cdot \vec{D}(y) \sigma_3, & F(-y) &= (\mathbb{1} \otimes \sigma_3) F(y) \sigma_3, & \Phi(-y) &= -\Phi(y),\end{aligned}\tag{5}$$

and charge operator is given by $Q = -q \otimes \sigma_3$.

The orbifold conditions and the reality conditions can be “solved” simultaneously using the relations

$$h = \begin{pmatrix} \phi_-^* & \phi_+ \\ -\phi_+^* & \phi_- \end{pmatrix}, \quad F = \begin{pmatrix} f_+ & -f_-^* \\ f_- & f_+^* \end{pmatrix}, \quad \chi = \begin{pmatrix} \lambda \\ -\lambda^{C_+} \end{pmatrix}, \quad \zeta = \begin{pmatrix} \psi \\ -\psi^{C_+} \end{pmatrix}, \quad \varepsilon = \begin{pmatrix} \eta \\ -\eta^{C_+} \end{pmatrix}.\tag{6}$$

The Dirac fermion λ can be decomposed into chiral projection on four dimensional Majorana spinors $\lambda_\pm$ as $\lambda = \lambda_{+L} - \lambda_{-R}$ and $\lambda^{C_+} = \lambda_{+R} + \lambda_{-L}$. The subscript $\pm$ refers to the eigenvalues under the parity operator $\sigma_3 i \gamma^5$. The right-handed states are not independent from the left-handed ones: $(\lambda_{-R})^{C_-} = \lambda_{-L}$, etc.. The supersymmetry parameter ε and the hyperino ζ can be represented by four dimensional Majorana spinors in a similar fashion. The parities of all fields in the model are given by the following table

$V :$	state	A_μ	A_5	Φ	$\lambda_{\pm L}$	$\lambda_{\pm R}$	D_3	$D_{1,2}$	$H :$	state	$\phi_\pm$	$\psi_{\pm L}$	$\psi_{\pm R}$	$f_\pm$
	parity	+	-	-	$\pm$	$\pm$	+	-		parity	$\pm$	$\pm$	$\pm$	$\pm$

The supersymmetry parameter η_+ corresponds to the 4 unbroken supersymmetry charges ($\eta_- = 0$). As observed in [11], the transformations of the odd and even fields reside in different multiplets with respect to this $N = \frac{1}{2}$ supersymmetry from the five dimensional point of view. For the even fields the transformation rules under this unbroken supersymmetry read

$$\begin{aligned}\delta A_\mu &= i \bar{\eta}_+ \gamma_\mu \lambda_+, & \delta \lambda_+ &= \frac{1}{4} F^{\mu\nu} \gamma_{\mu\nu} \eta_+ - \frac{1}{2} i \tilde{D}_3 \eta_+, & \delta \tilde{D}_3 &= \bar{\eta}_+ \gamma^\mu \partial_\mu \lambda_+, \\ \delta \phi_+ &= i \bar{\eta}_{+R} \psi_{+L}, & \delta \psi_{+L} &= -\gamma^\mu D_\mu \phi_+ \eta_{+R} - \tilde{f}_+ \eta_{+L}, \\ \delta \tilde{f}_+ &= \bar{\eta}_{+L} (i \gamma^\mu D_\mu \psi_{+L} + 2g_5 q \phi_+ \lambda_{+R}),\end{aligned}\tag{7}$$

but with “modified” auxiliary fields $\tilde{D}_3 = D_3 - \partial_y \Phi$ and $\tilde{f}_+ = f_+ - \partial_y \phi_-^*$. For odd fields similar transformations rules can be given. However, as the even fields (unlike the odd fields) do not vanish on the branes, they form $N = 1$ four dimensional supersymmetric multiplets on the branes: a vector multiplet $V| = (A_\mu, \lambda_+, \tilde{D}_3)$ and chiral multiplets $C_+| = (\phi_+, \psi_{+L}, \tilde{f}_+)$.

In addition to these multiplets that are induced from the vector and hyper multiplets in the bulk, we assume that there are an arbitrary number of chiral multiplets $C_0 = (\phi_0, \psi_{0L}, \tilde{f}_0)$ and $C_\pi = (\phi_\pi, \psi_{\pi L}, \tilde{f}_\pi)$ on the branes at $y = 0$ and $y = \pi R$, respectively. Their supersymmetry transformation rules are identical to those of $C_+|$. Using the standard four dimensional tensor calculus invariant Lagrangians can be obtained:

$$L_{branes} = \sum_{I=0,\pi} \delta(y - IR) \left\{ D_\mu \phi_I^\dagger D^\mu \phi_I + i \bar{\psi}_{IL} \not{D} \psi_{IL} + f_I^\dagger f_I + \right. \\ \left. + g_5 \phi_I^\dagger q_I \phi_I (-D_3 + \partial_y \Phi) - 2g_5 (\bar{\psi}_{IL} \lambda_{+R} q_I \phi_I + \text{h.c.}) \right\}. \quad (8)$$

3 Tadpoles

In the previous section the (unbroken) supersymmetry transformations for bulk vector (and hyper) multiplets have been discussed. Not D_3 but $\tilde{D}_3 = D_3 - \partial_y \Phi$ is the relevant auxiliary field for the remaining supersymmetry; this is true in particular on the branes. The one-loop FI-terms, due to brane chiral multiplets C_0 or C_π , are proportional to $\tilde{D}_3$; this is obvious as for them $\tilde{D}_3$ is the relevant auxiliary component. (The non-renormalization of the FI-terms in four dimensions has been proven in [15].) If supersymmetry is preserved at the one-loop level one would expect a similar result for the FI-tadpole due to hyper multiplets in the bulk: again this tadpole should be proportional to $\tilde{D}_3$. In ref. [2] it was shown that hyper multiplets may lead to a quadratically divergent FI-term, and in refs. [3, 4] it was argued that the counter term for D_3 is located at the branes. For completeness we redo a similar calculation here, but at the same time we show that also $\partial_y \Phi$ counter terms are needed on the boundaries, as one would expect from the supersymmetry analysis of the previous section. In ref. [1] the source of the divergence of Φ from bulk fields was identified to be the bulk fermions: the hyperinos.

Let us start with some more formal arguments to determine the generic structure of possible tadpoles. Notice that the Lagrangian (4) is quadratic in the hyper multiplet fields, hence the path integral over the hyper multiplets is formally trivial to compute and gives rise to determinants. Expanding to first order in Φ and $\vec{D}$ this gives us

$$i\Gamma_{eff} = \ln \int \mathcal{D}h \mathcal{D}\zeta e^{iS_H} \approx \text{tr}_\zeta \ln \left(\not{\partial} - i\frac{1}{2} \Phi Q \right) - \text{tr}_h \ln \left(i\Box \otimes \mathbb{1} + \frac{1}{2} i g_5 Q \otimes \vec{\sigma} \cdot \vec{D} \right) \\ \approx -g_5 \text{tr}_\zeta \left(\Box^{-1} i \not{\partial} \Phi Q \right) + \frac{1}{2} g_5 \text{tr}_h \left(\Box^{-1} Q \otimes \vec{\sigma} \cdot \vec{D} \right). \quad (9)$$

The zeroth order term cancels out because of supersymmetry. The traces over the hyperons and hyperinos, of course, have to be invariant under the boundary conditions (5). This shows that of the fermion and boson traces only $\text{tr}_\zeta \left(\Box^{-1} i \gamma^5 \partial_y \Phi Q \right)$ and $\text{tr}_h \left(\Box^{-1} Q D_3 \right)$ survive, respectively. Notice that this in particular shows, that the D_1, D_2 do not receive tadpole corrections. And that the field Φ will only appear with a derivative acting on it: $\partial_y \Phi$. Next, we compute those remaining traces using mode expansions.

As this calculation can be preformed using standard Feynman rules if we employ Dirac spinors and complex scalars instead of symplectic Majorana fermions and hyperons, we rewrite the relevant part of the hyper multiplet Lagrangian as

$$\begin{aligned}
L_H &\supset D_M \phi_+^\dagger D^M \phi_+ + D_M \phi_-^\dagger D^M \phi_- - g_5 D_3 (\phi_+^\dagger q \phi_+ - \phi_-^\dagger q \phi_-) \\
&\quad + i \bar{\psi} \gamma^M D_M \psi - g_5 \Phi \bar{\psi} q \psi, \\
L_{branes} &\supset \delta(y) \left(D_\mu \phi_0^\dagger D^\mu \phi_0 + g_5 (-D_3 + \partial_y \Phi) \phi_0^\dagger q_0 \phi_0 \right) \\
&\quad + \delta(y - \pi R) \left(D_\mu \phi_\pi^\dagger D^\mu \phi_\pi + g_5 (-D_3 + \partial_y \Phi) \phi_\pi^\dagger q_\pi \phi_\pi \right).
\end{aligned} \tag{10}$$

(This shows that the original hyper multiplet was normalized such that the standard normalization of complex bosons and Dirac fermions is obtained.)

The mode expansions, into even and odd state w.r.t. the parity, of the hyperons $\phi_\pm$, hyperinos $\psi_\pm$, the gauge scalar Φ and the auxiliary field D_3 read

$$\begin{aligned}
\phi_+(y) &= \eta_n \cos \frac{ny}{R} \phi_+^n, & \psi_{+L}(y) &= \eta_n \cos \frac{ny}{R} \psi_{+L}^n, & D_3(y) &= \eta_n \cos \frac{ny}{R} D_n, \\
\phi_-(y) &= \eta_n \sin \frac{ny}{R} \phi_-^n, & \psi_{-R}(y) &= \eta_n \sin \frac{ny}{R} \psi_{-R}^n, & \Phi(y) &= \eta_n \sin \frac{ny}{R} \Phi_n,
\end{aligned} \tag{11}$$


where, to take care of the different normalization of the zero mode, $\eta_0 = 1/\sqrt{\pi R}$ and $\eta_{n>0} = \sqrt{2/\pi R}$. In terms of these mode functions the propagators for the hyperons $\phi_\pm^n$ and the hyperinos $\psi_\pm^n$ become

$$\langle \phi_+^{n'\dagger} \phi_+^{n''} \rangle^{-1} = \frac{\delta_{n'n''}}{p_4^2 - n'^2/R^2}, \quad \langle \bar{\psi}_+^{n'\dagger} \psi_-^{n''} \rangle^{-1} = \frac{\delta_{n'n''}}{p_4^2 - n'^2/R^2} \begin{pmatrix} \not{p}_4 R & -\frac{n''}{R} L \\ -\frac{n''}{R} R & \not{p}_4 L \end{pmatrix}, \tag{12}$$

respectively. And the vertices of D_3 with $\phi_\pm$ and Φ with $\psi_\pm$ decomposed in terms of these modes give

$$\begin{aligned}
\langle D_n \phi_+^{n'\dagger} \phi_+^{n''} \rangle &= -g_5 q \frac{\pi R}{4} \eta_n \eta_{n'} \eta_{n''} \left(\delta_{n',n+n''} + \delta_{n'',n+n'} + \delta_{n,n'+n''} + \delta_{n+n'+n'',0} \right), \\
\langle D_n \phi_-^{n'\dagger} \phi_-^{n''} \rangle &= g_5 q \frac{\pi R}{4} \eta_n \eta_{n'} \eta_{n''} \left(\delta_{n',n+n''} + \delta_{n'',n+n'} - \delta_{n,n'+n''} - \delta_{n+n'+n'',0} \right), \\
\langle \Phi_n \bar{\psi}_{+L}^{n'\dagger} \psi_{-R}^{n''} \rangle &= g_5 q \frac{\pi R}{4} \eta_n \eta_{n'} \eta_{n''} \left(-\delta_{n',n+n''} + \delta_{n'',n+n'} + \delta_{n,n'+n''} - \delta_{n+n'+n'',0} \right), \\
\langle \Phi_n \bar{\psi}_{-R}^{n'\dagger} \psi_{+L}^{n''} \rangle &= g_5 q \frac{\pi R}{4} \eta_n \eta_{n'} \eta_{n''} \left(+\delta_{n',n+n''} - \delta_{n'',n+n'} + \delta_{n,n'+n''} - \delta_{n+n'+n'',0} \right),
\end{aligned} \tag{13}$$

respectively. The factor $\frac{\pi R}{4}$ arise by performing the integral over the interval $[0, \pi R]$. With these intermediate results the diagrams where the hyperons $\phi_\pm$ and the hyperinos ψ run around in the loop



can be calculated. By putting the propagators for the bosons and fermions together with the appropriate vertices, we find

$$\begin{aligned}\xi_{D_3} &= g_5 q \frac{\pi R}{4} \sum_{n,n',n''} \eta_n \eta_{n'} \eta_{n''} 2\delta_{n,n'+n''} \int \frac{d^4 p_4}{(2\pi)^4} \frac{\delta_{n',n''}}{p_4^2 - \frac{n'^2}{R^2}} D_n \\ \xi_\Phi &= (-2)g_5 q \frac{\pi R}{4} \sum_{n,n',n''} \eta_n \eta_{n'} \eta_{n''} 2\delta_{n,n'+n''} \int \frac{d^4 p_4}{(2\pi)^4} \frac{\delta_{n',n''}}{p_4^2 - \frac{n'^2}{R^2}} \left(-\frac{n''}{R}\right) \Phi_n.\end{aligned}\tag{14}$$

Some comments are in order here: the factor (-2) is due the trace over chiral fermions. In particular, we have used here that because of the chiral projections in (12) in the propagators only terms proportional to n''/R survive. By using the consequences of the Kronecker deltas, this can be represented as the derivate of Φ . Evaluating these expressions further, gives the result

$$\xi_{bulk} = \sum_n g_5 \text{tr}(q) \int \frac{d^4 p_4}{(2\pi)^4} \frac{1}{p_4^2 - n^2/R^2} \eta_{2n} \left(-D_{2n} + (\partial_y \Phi)_{2n}\right).\tag{15}$$


In this work we are only interested in the counter term structure of the theory which is determined by the divergent parts of (one-)loop results. Following [4] the divergent part of the remaining four dimensional integral is regulated using the cut-off Λ scheme

$$\int \frac{d^4 p_4}{(2\pi)^4} \frac{1}{p_4^2 - n^2/R^2} \Big|_{div} = \frac{1}{16\pi^2} \left(\Lambda^2 - \frac{n^2}{R^2} \ln \Lambda^2\right).\tag{16}$$

This shows that there is a leading quadratic divergence, which one would expect for Fayet-Iliopoulos tadpoles. But in addition there is also a sub-leading logarithmic divergence, that needs to be canceled by a counter terms as well. (Using dimensional regularization one can derive similar conclusions provided one introduces an additional infrared regulator to pick up the effects due to zero-modes.) In terms of a coordinate space representation of the divergent tadpoles, the y dependent FI-parameter then reads

$$\xi_{bulk}(y) = g_5 \frac{\text{tr}(q)}{2} \left(\frac{\Lambda^2}{16\pi^2} + \frac{\ln \Lambda^2}{16\pi^2} \frac{1}{4} \partial_y^2 \right) \left[\delta(y) + \delta(y - \pi R) \right].\tag{17}$$

We will see in later sections that the appearance of the double derivative of the delta-functions has important consequences. In addition to these bulk contributions, the diagrams



indicate that from the charged brane scalars we obtain the standard -4 -dimensional – FI-terms

$$\xi_{branes}(y) = g_5 \int \frac{d^4 p_4}{(2\pi)^4} \frac{1}{p_4^2} \sum_{I=0,\pi} \delta(y - IR) \text{tr}(q_I) = g_5 \frac{\Lambda^2}{16\pi^2} \sum_{I=0,\pi} \delta(y - IR) \text{tr}(q_I).\tag{18}$$

This together with the bulk FI-terms gives the full FI-parameter

$$\xi(y) = \xi_{bulk}(y) + \xi_{branes}(y).\tag{19}$$

4 Effective four dimensional potential

An analysis of the effective potential of this model is discussed next. With the notion “effective potential” is meant the part of the Lagrangian that can acquire Vacuum Expectation Values (VEVs) that do not break four dimensional Poincaré invariance. This implies that we allow for background solutions that are functions of y . As our analysis of this effective potential is essentially classical, we just assume that there are FI-terms on the two boundaries

$$L_{FI} = \left(-D_3 + \partial_y \Phi \right) \sum_{I=0,\pi} \left(\xi_I + \xi_I'' \partial_y^2 \right) \delta(y - IR), \quad (20)$$

with values ξ_0, ξ_0'' and ξ_π, ξ_π'' , respectively. The double derivative of the delta-functions was not considered in the classical FI-term analysis of refs. [6, 7]. If we assume that the FI-terms are dominated by the one-loop induced values we see that

$$\xi_I = g_5 \frac{\Lambda^2}{16\pi^2} \left(\frac{1}{2} \text{tr}(q) + \text{tr}(q_I) \right), \quad \xi_I'' = \frac{1}{4} g_5 \frac{\ln \Lambda^2}{16\pi^2} \frac{1}{2} \text{tr}(q). \quad (21)$$

Notice, in particular, that $\xi_0'' = \xi_\pi''$ when they are generated at the one-loop level. The FI-parameters ξ_0 and ξ_π receive contributions from both fields in the bulk and on the respective brane, while $\xi_0'' = \xi_\pi''$ only receives bulk contributions. Therefore the one-loop FI-term is completely absent, if the sum of the charges in the bulk and on both branes vanish separately.

By simply collecting the various relevant terms from the Lagrangians L_V , L_H , L_{brane} and L_{FI} an expression for the effective potential is obtained that is not manifestly semi-positive. However, by adding and subtracting the term $\frac{1}{2} \partial_y \Phi \text{tr}[h^\dagger Q h \sigma_3]$ and by doing an integration by parts on the subtracted one and using that the boundary terms are absent since Φ is odd, the effective potential takes the form

$$\begin{aligned} V_{4D} = \int_0^{\pi R} dy \quad & \left\{ \frac{1}{2} \left(\partial_y \Phi - \xi + \frac{1}{2} g_5 \text{tr}[h^\dagger Q h \sigma_3] + \delta(y) \phi_0^\dagger q_0 \phi_0 + \delta(y - \pi R) \phi_\pi^\dagger q_\pi \phi_\pi \right)^2 \right. \\ & - \frac{1}{2} \left(-D_3 - \xi + \frac{1}{2} g_5 \text{tr}[h^\dagger Q h \sigma_3] + \delta(y) \phi_0^\dagger q_0 \phi_0 + \delta(y - \pi R) \phi_\pi^\dagger q_\pi \phi_\pi \right)^2 \\ & + \frac{1}{8} g_5^2 \sum_{i=1,2} \left(\text{tr}[h^\dagger Q h \sigma_i] \right)^2 - \frac{1}{2} \sum_{i=1,2} \left(-D_i + \frac{1}{2} g_5 \text{tr}[h^\dagger Q h \sigma_i] \right)^2 + \\ & \left. + \frac{1}{2} \text{tr} \left[(\partial_y h^\dagger - i g_5 A_5 h^\dagger Q - g_5 \Phi \sigma_3 h^\dagger Q) (\partial_y h + i g_5 A_5 Q h - g_5 \Phi Q h \sigma_3) \right] \right\}. \quad (22) \end{aligned}$$

At first sight, it might seem that this expression is ill-defined because of the appearance of delta-functions squared. However, working out the squares shows that all (dangerous) delta-functions squared, in fact, drop out. Furthermore this expression is manifestly semi-positive: the negative squares that appear in the potential are zero using the equations of motions for D_3 and D_i .

Before we study supersymmetric background solutions in detail, let us say a few words about an important general property of background solutions on the boundaries. The vacuum field equations for $\phi_0^\dagger$ and $\phi_\pi^\dagger$ are trivial to determine:

$$q_0 \langle \phi_0 \rangle \langle -D_3 + \partial_y \Phi \rangle (0) = 0, \quad q_\pi \langle \phi_\pi \rangle \langle -D_3 + \partial_y \Phi \rangle (\pi R) = 0. \quad (23)$$

If none of the states in $\langle \phi_0 \rangle$ or $\langle \phi_\pi \rangle$ are chargeless, these equations imply that either supersymmetry is unbroken on the branes or the $U(1)$ symmetry is unbroken.

4.1 Supersymmetric background solutions

Supersymmetric vacuum solutions form a special class of all possible solutions. As usual their structure is easier to analyse than that of arbitrary (not necessarily supersymmetric) solutions. The equations that have to be satisfied in such a case are

$$\begin{aligned} D_3 = \partial_y \Phi &= \frac{1}{2} g_5 \text{tr}[h^\dagger Q h \sigma_3] + \delta(y) \phi_0^\dagger q_0 \phi_0 + \delta(y - \pi R) \phi_\pi^\dagger q_\pi \phi_\pi + \xi(y), \\ \text{tr}[h^\dagger Q h \sigma_i] &= 0, \quad \partial_y h + i g_5 A_5 Q h - g_5 \Phi Q h \sigma_3 = 0. \end{aligned} \quad (24)$$

For our subsequent discussion it is more convenient to leave the manifest 5 dimensional $N = 1$ supersymmetric notation, since it will turn out that the bosonic components, ϕ_+ and ϕ_- , have entirely different behavior. By substituting the solution (6) of the reality condition for the hyperon h , these relations become

$$D_3 = \partial_y \Phi = g_5 (\phi_+^\dagger q \phi_+ - \phi_-^\dagger q \phi_-) + \delta(y) g_5 \phi_0^\dagger q_0 \phi_0 + \delta(y - \pi R) g_5 \phi_\pi^\dagger q_\pi \phi_\pi + \xi(y), \quad (25)$$

$$\phi_+^T q \phi_- = 0, \quad \partial_y \phi_\pm \mp g_5 q \Phi \phi_\pm = 0. \quad (26)$$

In addition, A_5 has been put to zero. This can be achieved by a gauge transformation, since it vanishes on both boundaries already.

For a supersymmetric background solution these equations have to be satisfied simultaneously, and in addition the fields have to satisfy the appropriate boundary conditions. These equations show that four dimensional supersymmetry only requires that $D_3 - \partial_y \Phi$ vanishes, but not D_3 as one might expect. (This would be required if the five dimensional supersymmetry is to be preserved, but this is of course not possible while enforcing the orbifold conditions, in general). Non-vanishing odd fields are often not allowed because they are in conflict with the boundary conditions.

For the parity-odd bulk scalars ϕ_- the boundary condition implies that their VEVs are zero: $\langle \phi_- \rangle = 0$. This can be seen by the following argument: The solution in the bulk for ϕ_- is easily found. But since ϕ_- is odd, the integral over its derivative vanishes on $[0, \pi R]$. From this it follows that

$$0 = \int_0^{\pi R} dy \partial_y \langle \phi_- \rangle = \exp \left\{ -g_5 \int_0^{\pi R} dy \langle \Phi \rangle q \right\} \langle \phi_{0-} \rangle, \quad (27)$$

with $\langle \phi_{0-} \rangle$ the integration constant. Hence, unless the integral over $\langle \Phi \rangle$ diverges, the integration constant has to be zero. In particular, this shows that the equations involving ϕ_- in the supersymmetry conditions (26) are fulfilled trivially: $\langle \phi_{0-} \rangle = 0$.

The scalar of the gauge multiplet Φ is odd as well, it vanishes on the boundaries, and this leads to the integrability condition

$$0 = \int_0^{\pi R} dy \partial_y \Phi = \int_0^{\pi R} dy \langle \phi_{0+} \rangle^\dagger g_5 q \exp \left\{ 2g_5 \int_0^y dy \langle \Phi \rangle q \right\} \langle \phi_{0+} \rangle + \frac{1}{2} \sum_I \left(\langle \phi_I \rangle^\dagger g_5 q_I \langle \phi_I \rangle + \xi_I \right). \quad (28)$$

Here we have used that $\langle \phi_- \rangle = 0$. There is no contribution proportional to ξ_I'' since the first derivative of the delta function vanishes on the boundaries. Only the zero mode $D_{30} = \frac{1}{\pi R} \int_0^{\pi R} dy D_3$ of the auxiliary field D_3 is relevant for the question whether supersymmetry is broken due to gauge interactions (D-terms).

Let us analyze under what conditions this necessary (though in general not sufficient) requirement (28) for supersymmetric background is satisfied for the two distinct cases: broken or unbroken gauge symmetry. Suppose first that a background that does not spontaneously break the $U(1)$ gauge symmetry is possible. In that case all charged scalars (in the bulk and on the branes) have to vanish: $\langle\phi_+\rangle = 0$ and $\langle\phi_I\rangle = 0$. This is only possible if the sum of the brane FI-parameters vanishes:

$$\xi_{tot} = \xi_0 + \xi_\pi = 0 \quad \Rightarrow \quad \text{tr}(q) + \text{tr}(q_0) + \text{tr}(q_\pi) = 0. \quad (29)$$

This gauge invariant background is unique if all charges have the same sign. Otherwise, there can be many flat directions which spontaneously break the gauge symmetry. The relation of the sum of charges follow upon assuming that the FI-terms are induced at the one-loop level. When the FI-terms are generated by such quantum effects, it is not possible that all charges have the same sign since then the sum of the charges in (29) cannot be zero. Therefore, the one-loop FI-terms have flat directions that break gauge symmetry spontaneously.

If the sum of the FI-contributions $\xi_{tot} = \xi_0 + \xi_\pi$ is not equal zero, then supersymmetry can be maintained (i.e. eq. (28) is satisfied), provided that there is at least one (brane or bulk) field with a charge which has the sign opposite to ξ_{tot} . Hence in this case the gauge symmetry is always spontaneously broken. The gauge symmetry can then be broken by bulk or brane fields. Hence we see a similar situation then in 4 dimensions, if the (total integrated) FI-contribution is non-zero then either supersymmetry or gauge symmetry is spontaneously broken.

5 Localization of charged bulk fields

As we have observed in [1] the divergent FI-terms on the branes signal that the theory has a severe instability, that can lead to localization. The next topic we want to investigate in the paper is under what conditions charged bulk field can become localized and what the form of this localization is. In the first subsection, we answer when localization is certainly not possible, i.e. when the zero mode is constant. This is important because it gives us a criterion to decide what kind of bulk configuration are stable under quantum corrections. It occurs that the constant zero mode can be stable only in a special setup. Next, we investigate the possible shapes of the zero modes in the second subsection. The analysis that is performed here, only applies when we have a background that does not spontaneously break the gauge invariance. (The analysis can be extended to that case, but then not all even fields have zero modes due to the Higgs mechanism.) The main results of this analysis are summarized in tables 1 and 2. We end this section with a discussion of the massive part of the spectrum.

5.1 Conditions for constant zero mode

Because we only consider vacua that do not spontaneously break the gauge invariance, the relevant part Lagrangian which describes the bulk scalars is quadratic in them, it follows that the equation that determines the shape of the zero-mode is the same as the equation that determines the background solution. (Of course the zero mode has to be properly normalized.) As was argued, using eq. (27), the odd bulk fields can never have a non-trivial background solution. Hence there are no massless odd zero-modes.

The shape of the zero mode of a single even field⁴ ϕ_+ with charge q_b can formally be stated as

$$\partial_y \phi_{0+} - g_5 q_b \langle \Phi \rangle \phi_{0+} = 0 \quad \Rightarrow \quad \phi_{0+}(y) = \exp \left\{ g_5 q_b \int_0^y dy \langle \Phi \rangle \right\} \bar{\phi}_{0+}, \quad (30)$$

and is therefore entirely determined by $\langle \Phi \rangle$. The constants $\bar{\phi}_{0+}$ have to be chosen such that the zero mode is normalized as

$$\int_0^{\pi R} dy |\phi_{0+}(y)|^2 = 1. \quad (31)$$

From (30) it is clear that for $\langle \Phi \rangle(y) = 0$, the zero mode is constant over the 5th dimension.

Before investigating the properties of such localized zero modes further, we first turn to the question what the conditions are for the zero modes to be localized at all. In fact, it is easier first to answer when the zero mode does not get localized. Since $\langle \Phi \rangle$ must vanish for all y in that case, also its derivative vanishes. But then, since eq. (25) has to hold for all y , it follows that the conditions

$$\xi_I = 0, \quad \xi_I'' = 0, \quad (32)$$

have to be satisfied simultaneously. (The reason is that the delta functions, their derivatives and the zero modes are independent 'functions' (distributions) on the interval.) With the assumption that the FI-terms are only one-loop induced, these conditions given above become

$$\text{tr}(q_I) = 0, \quad \text{tr}(q) = 0. \quad (33)$$

The importance of these relations is that they indicate which configuration of supersymmetric bulk and brane fields are stable under quantum corrections. As we will see in the next subsections, if these conditions are not satisfied the description of the theory at the quantum level may be very different from the initial set-up, due to localization effects.

5.2 Shape of the zero mode

In this subsection we determine the profile of the scalar zero mode. As the model we consider is supersymmetric, it is to be expected that the chiral fermionic zero mode has the same shape over the extra dimension. This can also be seen directly from its equation of motion

$$(i\gamma^5 \partial_y - g_5 q_b \langle \Phi \rangle) \psi_{0+L} = 0 : \quad (34)$$

upon using that the eigenvalue of the chirality operator $i\gamma^5$ on the chiral state ψ_{0+L} is +1, this equation becomes identical to the scalar zero mode equation (30). Therefore, we now only focus on the scalar zero mode, knowing that all our conclusions apply to the chiral zero mode as well. We will make use of this fact when we discuss anomalies in section 6.3. At the end of this subsection we consider the limit $|\xi_0| \rightarrow \infty$ of our results, corresponding to taking the cut-off to infinity.

⁴Notice that we slightly abuse notation here since ϕ_+ denotes a vector of scalars in general, while now we refer to one component with charge q_b . By the subscript b we indicate that we really mean one component, and not the whole vector.

The extent of the localization of the zero mode (30) is determined by the solution of the equation for $\langle \Phi \rangle$ for a gauge invariant background ($\langle \phi_{0+} \rangle = 0$ and $\langle \phi_I \rangle = 0$) given by

$$\partial_y \langle \Phi \rangle = \left(\xi_0 + \xi_0'' \partial_y^2 \right) \delta(y) + \left(-\xi_0 + \xi_0'' \partial_y^2 \right) \delta(y - \pi R). \quad (35)$$

Here we have used that $\xi_0'' = \xi_\pi''$, and that only for $\xi_0 = -\xi_\pi$ the integrability condition (29) can be satisfied for a gauge and supersymmetric background. The solution of this equation is obtained straightforwardly

$$\langle \Phi \rangle(y) = \frac{1}{2} \xi_0 \operatorname{sgn}(y) + \xi_0'' \left(\delta'(y) + \delta'(y - \pi R) \right), \quad \operatorname{sgn}(y) = \begin{cases} 1 & 0 < y < \pi R, \\ 0 & y = 0, \pi R, \\ -1 & \pi R < y < 2\pi R. \end{cases} \quad (36)$$

And its integral

$$\int_0^y dy \langle \Phi \rangle(y) = \frac{1}{2} \xi_0 \left(\pi R - |y - \pi R| \right) + \xi_0'' \left(\delta(y) + \delta(y - \pi R) \right) \quad (37)$$

can be used to determine the shape of the zero modes in the presence of this special background. For this purpose we have to show explicitly what we obtain, when we formally insert the integral of Φ , eq. (37) into the form of the zero mode (30). The main problems here are: how to interpret the delta-functions in the exponential, and how to take the normalization condition into account.

To perform explicit calculations we need to regularize the delta-function in a suitable way.⁵ To this end the simple step-function is sufficient

$$\delta(y) \rightarrow \delta_\rho(y) = \begin{cases} \frac{1}{2\rho} & |y| < \rho, \\ 0 & |y| > \rho. \end{cases} \quad (38)$$

It reproduces the delta-function in the limit $\rho \rightarrow 0$. As this is a very simple prescription of the delta-function, the normalization constant $\bar{\phi}_{\rho+}$ of zero mode $\phi_{\rho+}(y) = \exp \left\{ g_5 q_b \int_0^y dy \langle \Phi \rangle \right\} \bar{\phi}_{\rho+}$ can be computed explicitly

$$\bar{\phi}_{\rho+}^{-2} = e^{g_5 q_b \xi_0'' / \rho} \left[\frac{e^{g_5 q_b \xi_0 \rho} - 1}{g_5 q_b \xi_0} + e^{g_5 q_b \xi_0 \pi R} \frac{1 - e^{-g_5 q_b \xi_0 \rho}}{g_5 q_b \xi_0} \right] + \frac{e^{g_5 q_b \xi_0 (\pi R - \rho)} - e^{g_5 q_b \xi_0 \rho}}{g_5 q_b \xi_0}. \quad (39)$$

Having determined the normalization factor with this regularization of the delta-function, next we investigate to what function the regularized zero mode $\phi_{\rho+}^2$ tends in the limit that $\rho \rightarrow 0$. Since the behavior of this limit strongly depends on the (relative) signs of various parameters, we now distinguish the following three cases: $\xi_0'' q_b > 0$, $\xi_0'' q_b < 0$, and $\xi_0'' = 0$. In the first case, where the charge q_b and the FI-parameter ξ_0'' have the same sign, the limit $\rho \rightarrow 0$ we obtain for the square of the zero mode is

$$\phi_{0+}^2(y) = \frac{2e^{g_5 q_b \xi_0 y}}{1 + e^{g_5 q_b \xi_0 \pi R}} [\delta(y) + \delta(y - \pi R)], \quad \xi_0'' q_b > 0. \quad (40)$$

⁵In a previous publication [1] we used a Gaussian regularization of the delta-function, to clarify the localization effects. To be able to perform all integrations explicitly we have chosen a different regularized form of the delta-function in this section.




Table 1: The two basic shapes (eqs. (40) and (41)) of the zero mode with charge q_b are displayed for a finite value of the cut-off Λ . (In table 2 the possible shapes are given in the infinite cut-off limit.) Delta function localizations, denoted by the arrows, happens if the signs of the FI-parameter ξ_0'' and charge q_b are the same. When the signs are opposite, the wave function falls off exponentially, and in addition vanishes at both branes. To which of the two branes the zero mode gets more localized is decided by the sign of $(\xi_0 - \xi_\pi)q_b$.

Hence the zero mode has the delta function support on the two fixed points, but the height at these two fixed points is not the same. In the second case, we find that the zero mode does not live on the boundaries of the interval

$$\phi_{0+}^2(y) = \frac{g_5 q_b \xi_0 e^{g_5 q_b \xi_0 y}}{e^{g_5 q_b \xi_0 \pi R} - 1} \begin{cases} 1 & 0 < y < \pi R, \\ 0 & y = 0, \pi R, \end{cases} \quad \xi_0'' q_b < 0. \quad (41)$$

This shows that the shape in this case is somewhat peculiar: the zero mode vanishes at both branes identically, while having an exponential behavior on the open interval $]0, \pi R[$. In table 1 we have schematically drawn both possible forms of localization. Notice that in both results the FI-parameter ξ_0'' does not appear anymore, since apart from its sign it can be completely absorbed in a rescaling of ρ which tends to zero anyway.

This rescaling is meaningless if ξ_0'' vanishes identically, and the zero mode then takes the form

$$\phi_{0+}^2(y) = \frac{g_5 q_b \xi_0 e^{g_5 q_b \xi_0 y}}{e^{g_5 q_b \xi_0 \pi R} - 1}, \quad \xi_0'' = 0. \quad (42)$$

A similar case has been studied before in refs. [7, 6].

Another intriguing possibility is when $\xi_0 = 0$. Eq. (40) then tells us that for $\xi_0'' q_b > 0$ the zero mode becomes a state that has two delta-function supports on both branes

$$\phi_{0+}^2(y) = \delta(y) + \delta(y - \pi R), \quad \xi_0 = 0, \quad \xi_0'' q_b > 0. \quad (43)$$

This is a dynamically generated state that has the property that it is localized on two (widely separated) branes. For the opposite case $\xi_0'' q_b < 0$ eq. (41) leads to a zero mode that is constant over the interior of the bulk but vanishes at both boundaries:

$$\phi_{0+}^2(y) = \begin{cases} 1 & 0 < y < \pi R, \\ 0 & y = 0, \pi R, \end{cases} \quad \xi_0 = 0, \quad \xi_0'' q_b < 0. \quad (44)$$

This state is not localized: it is constant over the bulk, but as it vanishes at the fixed points it cannot interact with brane fields.

When $\xi_0 \neq 0$ we can study what happens in the limit $|\xi_0| \rightarrow \infty$. For the case $\xi_0'' q_b \geq 0$ this limit leads to zero modes that are localized on one of the two branes

$$\phi_{0+}^2(y) = \begin{cases} 2\delta(y - \pi R) & \xi_0 q_b > 0, \\ 2\delta(y) & \xi_0 q_b < 0, \end{cases} \quad \xi_0'' q_b \geq 0. \quad (45)$$

The sign of $\xi_0 q_b$ decides on which of the two branes the state get localized. When $\xi_0'' q_b < 0$, we get an even more intriguing result: the zero mode is localized infinitely close to one of the fixed points; but on this fixed point itself the zero mode vanishes

$$\phi_{0+}^2(y) = \begin{cases} 2\delta(y - \pi R + 0) & \xi_0 q_b > 0, \\ 2\delta(y - 0) & \xi_0 q_b < 0, \end{cases} \quad \xi_0'' q_b < 0. \quad (46)$$

With the notation ± 0 we indicate that this zero mode is localized infinitely close, but not on a fixed point. From eqs. (45) and (46), one sees that the zero mode always gets localized at one of the two branes for very large $|\xi_0|$ irrespectively of the other FI-parameter ξ_0'' . However, for $\xi_0'' q_b < 0$ the zero mode does not actually end up on one of the two branes, but gets localized infinitely close to it.

In table 2 we have schematically drawn the various possibilities for the localization of the zero mode, depending on the charge of the bulk field and the signs of the brane FI-parameters. In these pictures we have taken the limit $|\xi_0| \rightarrow \infty$ when it is not zero.

The behavior of the bulk zero modes discussed in this subsection, can be viewed as a supersymmetric field theory analog of electrostatics. The system tries to distribute its free charges such that everywhere the net charge is zero. In this sense the configurations under investigation here are very similar to having two charged plates with some fluid in between which consists of charged particles as well. Also here the charge particles of the fluid distribute themselves, so as the net charge vanish everywhere if possible. This is because of the electric field generated by the differences of the charges of the two plates.

5.3 Mass spectrum

In the previous section we have seen that the scalar and chiral zero modes get localized on (one of) the branes. But these are only just the lightest modes of an infinite tower of states. Here we describe what happens to the masses of the other modes in the Kaluza-Klein tower for a finite FI-parameter ξ_0 , so that we clearly see what happens in the limit when this parameter tends to infinity.

As for the zero modes, we restrict our discussion of the mass spectrum to the scalars only. The masses for the fermions are identical, because their field equations can be “squared” to reproduce the boundary value problem of the scalars.

The spectrum of the bosonic Laplacian is given by the eigenvalue equation

$$\Delta\phi + \lambda\phi = 0, \quad \Delta = \partial_y^2 - g_5 q_b \left(\partial_y \langle \Phi \rangle + g_5 q_b \langle \Phi \rangle^2 \right), \quad (47)$$

where we concentrate on the case where $\langle \Phi \rangle$ is given by eq. (36), with the appropriate boundary conditions of the even and odd scalar fields imposed. As usual this ensures that the spectrum of

$$\xi_0'' \sim \ln(\Lambda^2) \text{tr}(q), \quad \xi_0 \sim \Lambda^2 \left(\text{tr}(q_0) + \frac{1}{2} \text{tr}(q) \right), \quad \text{SUSY} + U(1) \Rightarrow \xi_0 = -\xi_\pi$$

$$\xi_\pi \sim \Lambda^2 \left(\text{tr}(q_\pi) + \frac{1}{2} \text{tr}(q) \right),$$





$q_b \neq 0$	$\xi_0'' q_b \geq 0$	$\xi_0'' q_b < 0$
$\xi_0 q_b < 0$ ($\xi_\pi q_b > 0$)		
$\xi_\pi q_b < 0$ ($\xi_0 q_b > 0$)		
$\xi_0 = \xi_\pi = 0$ $\xi_0'' \neq 0$		

Table 2: This table schematically displays the different shapes of a (bulk) zero mode with charge $q_b \neq 0$, summarizing eqs. (45), (46), (43), and (44) with the cut-off $\Lambda \rightarrow \infty$. This mode localizes at the brane with the sign of its FI-parameter (ξ_0, ξ_π) opposite to the charge of the zero mode. To allow for a supersymmetric vacuum solution that preserves the $U(1)$ symmetry, these two FI-parameters add up to zero. Assuming that the one-loop FI-terms dominate, these FI-parameters are proportional to the sum of charges at the branes, and the third FI-parameter ξ_0'' is proportional to the sum of charges in the bulk. If the product of q_b and ξ_0'' is negative the zero mode localizes infinitely close to but not at the brane, otherwise it localizes exactly on the brane. Vanishing FI-terms ξ_0, ξ_π is special: if $\xi_0'' q_b > 0$ the zero mode splits up into two delta functions on both branes, while for $\xi_0'' q_b < 0$ the zero mode is constant over the interior of the bulk but vanishes at its boundaries. When all FI-parameters vanish, of course, there is no localization effect.

eigenvalues λ is discrete. To avoid having to deal with the second derivative of the delta-function explicitly, we perform the substitution $\phi = \exp \{g_5 q_b \int_0^y dy \langle \Phi \rangle\} \tilde{\phi}$ so as to obtain the boundary value problem

$$\tilde{\phi}'' + 2g_5 q_b \langle \Phi \rangle \tilde{\phi}' + \lambda \tilde{\phi} = 0, \quad \tilde{\phi}'(0) = \tilde{\phi}'(\pi R) = 0, \quad (48)$$

for the even fields. (For odd fields the boundary conditions are $\tilde{\phi}(0) = \tilde{\phi}(\pi R) = 0$.)

As this equation still involves a first derivative on the delta function, some appropriate regularization is needed. In appendix B such a regulator is provided and the algebraic equation that determines the eigenvalues is obtained. In the limit where this regulator is removed, the spectrum for various signs of the FI-parameters and even and odd fields becomes identical, and completely independent of the FI-parameter ξ_0'' . The mass spectrum reads

$$\lambda_n = \frac{1}{4}(g_5 q_b \xi_0)^2 + \frac{n^2}{R^2}, \quad n \in \mathbb{N}, \quad (49)$$

see (B.8). Clearly, in the limit $|\xi_0| \rightarrow \infty$ (i.e. when the cut-off Λ becomes extremely large) this shows that all non-zero modes become infinitely heavy, and that all these states should decouple from the theory. However, there might be some subtleties due to non-decoupling effects as we will see in our analysis of anomalies.

6 Anomalies

In this section we turn to the important discussion of anomalies in orbifold models. The first subsection is devoted to the analysis of a specific type of global anomaly for theories on $\mathcal{M}^4 \times S^1$: the parity anomaly. The importance of global anomalies has been studied extensively in the past [16, 17]. One of the first works on this was [18], which showed that a theory with an odd number of $SU(2)$ doublets is inconsistent. We will see that the parity anomaly leads to a similar conclusion for MSSM-like orbifold models with an odd number of $SU(2)$ doublets or $SU(3)$ triplets in the bulk. The central observation we make in this section is that before defining an orbifold model one should check whether the parity symmetry is a quantum symmetry because otherwise it may be problematic to divide it out to construct the orbifold.

The next section studies the local gauge anomalies that can arise in orbifold models. Using topological arguments it is shown that the gauge anomalies appear only on the boundaries. Conditions for consistency of an orbifold theory with fermions is given. In the last subsection we discuss to what extent the localization effects can change the discussion of anomalies.

6.1 Parity anomaly on S^1

In this subsection we point out that some orbifold models containing fermions are ill-defined. To explain what the problem is, we need to take one step back and consider a fermionic field theory on a circle in 5 dimensions coupled to a gauge field A_M associated to a Lie group G . On the circle this theory is classically invariant under the parity transformation

$$y \rightarrow -y, \quad A_\mu(y) \rightarrow A_\mu(-y), \quad A_5(y) \rightarrow -A_5(-y), \quad \psi(y) \rightarrow i\gamma^5 \psi(-y). \quad (50)$$

In the construction of the orbifold $S^1/\mathbb{Z}_2$ this symmetry is divided out.

But this symmetry can be anomalous as was observed in ref. [19], and can lead to a sign ambiguity in the fermionic determinant. This anomaly can be canceled by adding the parity anomaly counter term

$$\Gamma_{PAC}(A) = -\pi i \int_{\mathcal{M}^4 \times S^1} \Omega_5(A), \quad (51)$$

with $\Omega_5(A)$ the Chern-Simons 5-form defined in eq. (C.2) of appendix C. (There we briefly recall some standard facts about anomalies and characteristic classes to set up the notation used in the section.) The effective (Euclidean) quantum action $\Gamma(A)$ with the fermions integrated out is defined only modulo $2\pi i$, see eq. (C.3). Therefore, if the difference $\Delta\Gamma_{PAC}(A)$ between the parity transformed and the counter term (51) satisfies

$$\Delta\Gamma_{PAC}(A) = 2\pi i \int_{\mathcal{M}^4 \times S^1} \Omega_5(A) = 0 \text{ mod } 2\pi i, \quad (52)$$

the model does not have a parity anomaly. Since the APS η -invariant [20] $\eta = 2 \int \Omega_5(A) + \text{const.}$ is an integer, it follows that if $\int \Omega_5(A)$ is an even multiple of $\int \Omega_{5|F}(A)$ computed with the trace of the fundamental representation, there is no parity anomaly. As this Chern-Simons form is proportional to the symmetric trace $D_{ijk} = \text{tr}(T_i\{T_j, T_k\})$, standard four dimensional anomaly cancelation arguments can be used to argue when this counter term is just zero.

It is not hard to show that the parity counter term is invariant under infinitesimal gauge transformations, using similar arguments as in (C.6). However, this counter term Γ_{PAC} gives back a sign ambiguity by a large gauge transformation g if

$$\Gamma_{PAC}(^g A) \neq 0 \text{ mod } 2\pi i. \quad (53)$$

If this happens the counter term Γ_{PAC} cannot be added since then gauge invariance is lost. In appendix D we analyse, using homotopy groups, which parity anomaly counter terms are potentially not gauge invariant. We find that if the gauge group contains $SU(n)$ (with $n \geq 3$) or $USp(2n)$ factors, or is a product of at least one $U(1)$ and some simple groups, gauge invariance of the parity counter term depends on the precise choice of matter representation, otherwise gauge invariance is automatic. (Parity counter term for just $U(1)$ can also violate gauge symmetry if gravitational interactions are also taken into account.)

The parity anomaly for a field theory on the circle S^1 is not a problem, it just means that the parity invariance is broken at the quantum level. However, if one wants to define a field theory on the orbifold $S^1/\mathbb{Z}_2$, it is of course crucial that this parity symmetry is an exact symmetry of the quantum theory, else it makes no sense to mod it out!

Therefore, if the fermionic determinant on S^1 is not parity invariant, the parity counter term (51) has to be added. But then it is crucial that gauge invariance in the sense of eq. (53) is not lost! This reintroduces a sign-ambiguity in the path integral description of the theory with fermions and gauge fields. It is important to note here that the orbifolding condition does not project out those large $U(1)$ -gauge transformations under which the parity counter term is not invariant. For example, they can be maps of $g : S^1 \rightarrow U(1)$ that are topologically equivalent to the form $g(y) = e^{iy/(2\pi R)}$ which has non-trivial winding on $U(1)$. Hence the parity changes the orientation of the winding but does not undo it. If it is not possible to save the parity symmetry on S^1 using a gauge invariant counter term, then one can question whether the model with the parity divided out at the classical level, gives rise to a consistent quantum field theory.

We now investigate which MSSM-like models with fields in the bulk and/or on the branes do not suffer from the parity anomaly. According to the analysis above, we have to check that all possible parity anomaly counter terms appear as even multiples of those for the fundamental representations. This gives the following constraints for the bulk multiplets:

$$\left. \begin{array}{ll} U(1)^3 & 6n_Q q_Q^3 + 2n_L q_L^3 + 3n_{u^*} q_{u^*}^3 + 3n_{d^*} q_{d^*}^3 + n_{e^*} q_{e^*}^3 + 2n_{H_1} q_{H_1}^3 + 2n_{H_2} q_{H_2}^3 \\ SU(3)^3 & 2n_Q + n_{u^*} + n_{d^*} \\ U(1) \times SU(2)^2 & 3n_Q q_Q + n_L q_L + n_{H_1} q_{H_1} + n_{H_2} q_{H_2} \\ U(1) \times SU(3)^2 & 2n_Q q_Q + n_{u^*} q_{u^*} + n_{d^*} q_{d^*} \\ U(1) \times \text{grav} & 6n_Q q_Q + 2n_L q_L + 3n_{u^*} q_{u^*} + 3n_{d^*} q_{d^*} + n_{e^*} q_{e^*} + 2n_{H_1} q_{H_1} + 2n_{H_2} q_{H_2} \end{array} \right\} = 0 \text{ mod } 2, \quad (54)$$

where q_X is the hypercharge of multiplet X in units of the hypercharge of the quark doublet Q and n_X is the number of such multiplets present in the bulk. Three of the above conditions are fulfilled automatically for arbitrary n_X because q_{u^*} , q_{d^*} and q_{e^*} are even. Only the second and third relations give the non trivial restrictions:

$$\begin{aligned} n_{u^*} + n_{d^*} &= 0 \text{ mod } 2, \\ n_Q + n_L + n_{H_1} + n_{H_2} &= 0 \text{ mod } 2. \end{aligned} \quad (55)$$

So we conclude that in MSSM-like models the parity anomaly appears when an odd number of $SU(2)$ doublets (quark, lepton or Higgs) or an odd number of right-handed quarks live in the bulk. Hence we need a parity counter term that involves $U(1)$ and $SU(2)$, and $SU(3)$ gauge field interactions. However, this is precisely of the form of not necessarily gauge invariant parity counter terms as explained in appendix D. Indeed for an odd number of $SU(2)$ doublets or an odd number of $SU(3)$ triplets of the MSSM the conclusion is that the parity counter term is not gauge invariant.

Let us close this subsection by making some general comments clarifying the meaning and further applications of the results presented here. First of all an orbifold theory is a special theory on an interval in which, apart from the Dirichlet or Neumann boundary conditions for the fields, there is a symmetry that forbids many operators. By doubling the space this symmetry can be made explicit and is identified with the $\mathbb{Z}_2$ parity. Therefore the parity anomaly does not forbid a quantum field theory on an interval with appropriate boundary conditions, but it implies that many additional terms may appear in the Lagrangian as there is no (quantum) symmetry which forbids them.

Secondly, we notice that the parity anomaly may have (important) consequences when the mod out of the parity symmetry is used to break the gauge symmetry [21] (which may have interesting applications like solving the doublet-triplet splitting problem [22]). Here it is important to take into account the distinction between inner and outer automorphism gauge symmetry breaking (see e.g. [23]). For inner automorphisms we see, that the parity anomaly can arise under the same conditions as discussed in the subsection and appendix D (since with $A_\mu(-y) = P A_\mu(y) P^{-1}$, etc., the gauge group element P that squares to unity, drops out of the Chern-Simons term (52)). The parity anomaly may not arise when outer automorphisms are used to break the gauge group: for example, under complex conjugation of the generators of the gauge group $T_a \rightarrow (-T_a)^T$, the parity counter term is invariant.

6.2 Anomaly free models on $S^1/\mathbb{Z}_2$

In the recent literature [4, 8, 9, 10, 24] there has been a lot of attention to the discussion on the possible forms of local gauge anomalies on orbifolds. On the orbifold $S^1/\mathbb{Z}_2$, the standard argument that there cannot be anomalies as there are no chiral fermions does not hold because on the four dimensional boundaries chiral projection may appear. In this section we start with the possible gauge anomaly due to a five dimensional Dirac fermion coupled to a gauge field A_M on the orbifold $S^1/\mathbb{Z}_2$. After that we consider localization effects, additional chiral fermions on the boundaries, and discuss possible Chern-Simons counter terms.

Following Horava-Witten [12] the form of the anomaly for Dirac fermion ψ on $\mathcal{M}_5 = \mathcal{M}^4 \times S^1/\mathbb{Z}_2$, that satisfies $\psi(-y) = i\gamma^5\psi(y)$, can be determined as follows. First of all the anomaly has to be mathematically consistent, this means that the anomaly has to satisfy the Wess-Zumino consistency condition [25]. Since the solution $\Omega_4^1(A, \Lambda)$ to this condition is unique up to a normalization factor, the effective action $\Gamma(A)$, obtained by integrating the fermion out, transforms as

$$\delta_\Lambda \Gamma(A) = \pi i \sum_{I=0,\pi} \tilde{n}_I \int_{\mathcal{M}^4} \Omega_4^1(A; \Lambda) \Big|_{IR}, \quad e^{-\Gamma(A)} = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S}, \quad (56)$$

with only the constants $\tilde{n}_I$ left to be determined. (One could have added $\int_{\mathcal{M}_5} d\Omega_4^1(A, \Lambda)$, but this can be rewritten as two boundary terms, already considered in (56).) This argument shows that the anomaly can only appear on the 4 dimensional boundaries. Since the projections on both fixed points are the same, it implies that $\tilde{n}_0 = \tilde{n}_\pi$. By restricting both the gauge parameter and the fields to four dimensions, the standard result for a chiral fermion in 4 dimensions should be obtained: eq. (C.5) of the appendix C. This then fixes the normalization completely $\tilde{n}_0 = \tilde{n}_\pi = 1$.

Writing the anomaly as an integral over the five dimensional space $\mathcal{M}_5$

$$\delta_\Lambda \Gamma(A) = \pi i \int_{\mathcal{M}_5} (\delta(y) + \delta(y - \pi R)) \Omega_4^1(A; \Lambda) dy, \quad (57)$$

the same form of the anomaly is obtained as the one found by [8] using a perturbative calculation. It was observed there that the result was independent of the shape of the modes used to perform the calculation, provided of course, that the set of modes is complete. This conclusion is also reached from the reasoning given here: we see clearly the topological origin of this anomaly as it is defined in terms of the characteristic classes.

Next, we include the possibility of having chiral fermions on the boundaries. Under a gauge transformation the effective actions $\Gamma_I(A)$ of the fermions on the branes $I = 0, \pi$ can have anomalies

$$\delta_\Lambda \Gamma_I(A) = 2\pi i \int_{\mathcal{M}^4} \Omega_{4|I}^1(A; \Lambda) \Big|_{IR}, \quad e^{-\Gamma_I(A)} = \int \mathcal{D}\psi_I \mathcal{D}\bar{\psi}_I e^{-S_I}, \quad (58)$$

with $|I$ we indicate that the trace which is implicit in the form $\Omega_{4|I}^1$ is taken over the chiral fermions living on brane I only.

In order to obtain a consistent model, we may need to introduce possible Chern-Simons terms. The uniqueness of the descent equation (C.2) tells us that there is only one possible five dimensional Chern-Simons action $\Gamma_{CS}(A)$ available up to normalization

$$\Gamma_{CS}(A) = N_{CS} \pi i \int_{\mathcal{M}_5} \Omega_{5|F}(A), \quad (59)$$

where the internal trace is taken over the fundamental representation (F) to fix the normalization N_{CS} uniquely. The gauge transformation (C.1) of the Chern-Simons term reads

$$\delta_\Lambda \Gamma_{CS}(A) = N_{CS} \pi i \int_{\mathcal{M}_5} d\Omega_{4|F}^1(A; \Lambda) = N_{CS} \pi i \int_{\mathcal{M}_5} (-\delta(y) + \delta(y - \pi R)) \Omega_{4|F}^1(A; \Lambda) dy. \quad (60)$$

The minus sign in front of $\delta(y)$ appears because of changing the induced orientation, due to Stoke's theorem, to standard one.

A model in five dimensions on the orbifold $S^1/\mathbb{Z}_2$ with both bulk and brane fermions can be made locally consistent, if one can find a normalization N_{CS} of the Chern-Simons term, such that the total effective action

$$\Gamma_{tot}(A) = \Gamma(A) + \Gamma_0(A) + \Gamma_\pi(A) + \Gamma_{CS}(A) \quad (61)$$

is gauge invariant under local (in general five dimensional) gauge transformations. By normalizing the bulk and brane fermion anomalies (56) and (58) w.r.t. the fundament representation (F),

$$\begin{aligned} \delta_\Lambda \Gamma(A) &= N \pi i \int_{\mathcal{M}_5} (\delta(y) + \delta(y - \pi R)) \Omega_{4|F}^1(A; \Lambda) dy, \\ \delta_\Lambda \Gamma_I(A) &= 2N_I \pi i \int_{\mathcal{M}_5} \delta(y - IR) \Omega_{4|F}^1(A; \Lambda) dy, \end{aligned} \quad (62)$$

similarly to the Chern-Simons term, one can determine N_{CS} , and obtain a consistency condition

$$N_{CS} = N_0 - N_\pi, \quad N + N_0 + N_\pi = 0. \quad (63)$$

Notice that the consistency requirement takes the form of a sum rule and is determined by the fermionic content of the bulk and branes only. Furthermore, we see that a Chern-Simons term is required only if the anomalies due to both branes are not equal. It should be stressed that the results of this subsection depend on properties of the fermionic content of the model, therefore the requirements should hold in both supersymmetric and non-supersymmetric models on $S^1/\mathbb{Z}_2$.⁶

It is easy to apply these formulae to the case of possible pure $U(1)$ gauge and mixed $U(1)$ -gravitational anomalies, we find the anomaly cancelation conditions

$$\text{tr}(q^3) + \text{tr}(q_0^3) + \text{tr}(q_\pi^3) = 0, \quad \text{tr}(q) + \text{tr}(q_0) + \text{tr}(q_\pi) = 0. \quad (64)$$

Observe that these relations are exactly the expected ones by looking only at the low-energy spectrum determined by the zero modes. And the coefficients for the pure $U(1)$ and mixed Chern-Simons counter terms are given by

$$N_{CS}(U^3(1)) \sim \text{tr}(q_0^3) - \text{tr}(q_\pi^3), \quad N_{CS}(\text{mixed}) \sim \text{tr}(q_0) - \text{tr}(q_\pi), \quad (65)$$

up to normalization to the minimal charge.

⁶ A similar analysis can be preformed for the orbifold $S^1/\mathbb{Z}_2 \times \mathbb{Z}_2'$. In this case the chiral projection on one brane can be opposite to that on the other brane. Then one finds $\tilde{n}_0 = -\tilde{n}_\pi = 1$ in (56). By denoting the normalization of the anomaly in both cases by $N_\pm$, and following the same arguments, we find a consistency requirement: $N_+ + N_0 + N_\pi = 0$, involving matter with chiral zero modes only, while the normalization of the Chern-Simons term is given by $N_{CS} = N_- + N_0 - N_\pi$.


6.3 Gauge anomalies on $S^1/\mathbb{Z}_2$ and localization

The analysis of the previous subsection was performed for a quite general 5 dimensional model on the interval $S^1/\mathbb{Z}_2$ with fermions and gauge fields. We now return to a much more restrictive situation of a supersymmetric model.

Because the result for the anomaly is independent of which complete set of modes is used [8] (but of course with the same boundary conditions), it follows that the localization effects can not change the form of the anomaly. However, this does not mean that the localization does not change our description of the situation involving anomalies. When localization of the zero modes happen and the cut-off is taken to be very large, all modes except the zero mode get infinitely heavy, see section 5.3. Therefore for many practical purposes they have completely decoupled from the theory. Let us introduce some notation to describe the situation: Let $\Gamma^0(A)$ denote the effective action after integrating out the (localized) chiral zero mode (while neglecting the massive modes) and $\Gamma^M(A)$ the effective action obtained by integrating out the massive fermionic states (and dropping the massless one). Since the (localized) chiral zero mode and the massive fermion modes are independent in the path integral, it follows that the bulk effective action may be written as $\Gamma(A) = \Gamma^0(A) + \Gamma^M(A)$. From this equation, one can define the “massive” anomaly due to the (infinitely) heavy modes by

$$\delta_\Lambda \Gamma^M(A) = \delta_\Lambda \Gamma(A) - \delta_\Lambda \Gamma^0(A). \quad (66)$$


As the first term is given by (56), and the second we know since it represents the anomaly due to a localized state, the “massive” anomaly is given. As an example, let us give a graphical representation of this equation. Consider the case that the zero modes get localized at brane 0, this leads to the identity



$$\text{“massive” anomaly} = \text{bulk anomaly} - \text{localized zero mode anomaly}. \quad (67)$$

This “massive” anomaly has the same structure as the graphical representation of the gauge transformation of the Chern-Simons term (60). In a consistent model, one would expect that this “massive” anomaly is canceled by other high-energy physics effects like five dimensional Chern-Simons terms (see e.g. [24]), like it is done here when the zero mode splits up to exist on both branes only.

Let us consider a simple situation, that we have considered in our previous work [1]: a model with one chiral multiplet on brane 0 and a hyper multiplet of the opposite charge in the bulk. It is not hard to check that this model is anomaly free but needs ($N_{CS} = -1$) a $U(1)$ gauge Chern-Simons term (neglecting gravitational interaction here for simplicity) because the anomalies on both branes are not equal. Pictorially the anomaly cancelation in this model can be drawn as



$$\text{brane anomaly} + \text{bulk anomaly} - \text{Chern-Simons variation} = 0. \quad (68)$$

Applying our analysis of section 5 to this model, we infer the bulk field get localized at brane 0. Hence we may use the equality (67) to eliminate the gauge variation of the Chern-Simons term. The bulk anomaly contribution cancels out and we are left with local anomaly cancellation on the 0 brane.

$$\begin{array}{c} \text{brane} \\ \text{anomaly} \end{array} \quad \begin{array}{c} \text{localized zero} \\ \text{mode anomaly} \end{array} = 0. \quad (69)$$

After the localization the Chern-Simons term is absent, canceled so to say, against the anomaly due to the heavy modes, and the anomaly cancellation involves zero modes only.

7 Conclusions

In this work we have set out to investigate two aspects of supersymmetric theory in 5 dimensions compactified on the orbifold $S^1/\mathbb{Z}_2$ which may lead to a deeper understanding of structure and properties of such models: consistency requirements due to anomalies and instabilities due to Fayet-Iliopoulos terms.

After a review of supersymmetric theories in 5 dimensions with boundaries, we performed a one-loop calculation of the Fayet-Iliopoulos terms at the branes generated by brane and bulk fields. We showed that the bulk fermions are responsible for the generation of the tadpole for $\partial_y \Phi$, which is required to appear together with the auxiliary field $-D_3$ because of the unbroken supersymmetry.

A substantial part of the paper was devoted to the study of localization effects caused by (quadratically and logarithmically) divergent Fayet-Iliopoulos terms due to bulk and brane fields. We have studied the localization of zero modes in the presence of supersymmetric and $U(1)$ gauge symmetry preserving vacuum solutions. We have identified three different types of possible localized zero modes when the cut-off is taken very large: localization at a brane, localization infinitely close to (though not on) a brane, and simultaneous localization at both boundaries. The second form of localization leads to the situation that some fields live at the same place but any direct interaction between them is forbidden. Third type of localization leads to states that live and interact on two branes which may have arbitrary large distance between them.

It should be stressed that the localization discussed in this paper is typically of the delta-like type. The reason for this is the δ'' -contribution in the FI-tadpoles. Only the second kind of localization (close to a brane) may have some finite width which is controlled by the cut-off scale Λ (so this width may be substantial only when the cut-off scale is very low or when the tree level FI-term is fine-tuned against the radiatively generated one). This means that the localization effect is important, not only for models with constant zero modes of bulk fields, but also for models (for example [7, 26, 27]) in which the zero modes have more complicated shapes.

Another source of quantum instabilities, anomalies, were investigated in the remainder of the paper. First, we discussed a possible difficulty in defining the fermionic model on this orbifold at the quantum level, if on S^1 there is an anomaly in the $\mathbb{Z}_2$ parity symmetry. We derived sufficient requirements on the fermionic spectrum of theory, so as to avoid these complications.

Then we moved to local gauge anomalies on the orbifold $S^1/\mathbb{Z}_2$. Using the uniqueness of the solutions of the Wess-Zumino consistency condition for anomalies, the structure of anomalies due to bulk and brane fields could be easily determined. The role of possible Chern-Simons terms in anomaly cancellation was addressed: we showed that demanding 5 dimensional local gauge invariance leads to consistency conditions which are independent of the presences of Chern-Simons interactions. These conditions are nothing but the familiar 4 dimensional anomaly cancellation requirements for the zero modes of the theory. The normalization of the Chern-Simons terms, on the other hand, is determined by the imbalance of gauge anomalies due to the chiral fermions on the two branes. In addition, we showed, that although the localization of bulk zero modes to branes does not change the consistency requirement on the theory (since this only involve the zero modes), the remaining (infinitely) heavy modes may cancel (parts of) the Chern-Simons terms.

Acknowledgments

We would like to thank H.M. Lee, J.E. Kim and M. Walter for useful discussions.

Work supported in part by the European Community's Human Potential Programme under contracts HPRN-CT-2000-00131 Quantum Spacetime, HPRN-CT-2000-00148 Physics Across the Present Energy Frontier and HPRN-CT-2000-00152 Supersymmetry and the Early Universe. SGN was supported by priority grant 1096 of the Deutsche Forschungsgemeinschaft. MO was partially supported by the Polish KBN grant 2 P03B 052 16.

A Notations and Conventions

The metric has the signature $\eta_{MN} = (+1, -1, -1, -1, -1)$, with $M, N = 0, \dots, 4$. The Clifford algebra in five dimensions can be characterized by the following relations

$$\{\gamma^M, \gamma^N\} = 2\eta^{MN}, \quad \gamma^{M\dagger} = \gamma^0 \gamma^M \gamma^0, \quad C_+^{-1} \gamma^M C_+ = \gamma^{MT}, \quad C_+^T = -C_+, \quad C_+^\dagger = C_+^{-1}. \quad (\text{A.1})$$

Dirac and charge conjugation are given by

$$\bar{\psi} = \psi^\dagger \gamma^0, \quad \psi^{C+} = C_+ \bar{\psi}^T, \quad (\text{A.2})$$

respectively.

The symplectic structure of $N = 1$ in five dimensions can be made manifest using symplectic (Majorana) reality conditions

$$\varepsilon = \epsilon \varepsilon^{C+}, \quad \chi = \epsilon \chi^{C+}, \quad \zeta = \rho \zeta^{C+}, \quad h = \rho h^* \epsilon^{-1}, \quad (\text{A.3})$$

for the supersymmetry parameters ε , the gauginos $\chi = (\chi^\alpha)$, the hyperinos $\zeta = (\zeta^a)$ and the hyperons $h = (h^{a\alpha})$, with $\alpha = 1, 2$ and $a = 1, \dots, 2n$ for n hyper multiplets, respectively. The chirality operator $i\gamma^5$ and the matrices ϵ and ρ can be represented as

$$i\gamma^5 = \begin{pmatrix} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{pmatrix}, \quad \epsilon = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \rho = \mathbb{1} \otimes \epsilon = \begin{pmatrix} 0 & \mathbb{1}_n \\ -\mathbb{1}_n & 0 \end{pmatrix}. \quad (\text{A.4})$$

The five and four dimensional charge conjugation matrices are related via $C_+ = i\gamma^5 C_-$.

The Pauli-matrices, denoted by $\vec{\sigma} = (\sigma^A)$, carry the same indices as the auxiliary fields D^A of the vector multiplet. The different indices are suppressed as much as possible throughout this work.

B Scalar spectrum

In this appendix we compute the spectrum of the boundary value problem (48) for even fields in detail. Since the calculation for the odd fields is completely analogous, we conclude this appendix by simply quoting the results in that case.

The spectrum of the bosonic Laplacian is given by the eigenvalue equation

$$\Delta\phi + \lambda\phi = 0, \quad \Delta = \partial_y^2 - g_5 q_b \left(\partial_y \langle \Phi \rangle + g_5 q_b \langle \Phi \rangle^2 \right), \quad (\text{B.1})$$

where we concentrate on the case where $\langle \Phi \rangle$ is given by eq. (36). Taking ϕ to be even implies that on the interval $[0, \pi R]$ it has to satisfy the boundary conditions $\phi'(0) = \phi'(\pi R) = 0$. As usual this ensures that the spectrum of eigenvalues λ is discrete.

To avoid having to deal with the square of the first derivative of delta-functions explicitly, we perform the substitution $\phi = \exp \left\{ g_5 q_b \int_0^y dy \langle \Phi \rangle \right\} \tilde{\phi}$ so as to obtain the boundary value problem

$$\tilde{\phi}'' + 2g_5 q_b \langle \Phi \rangle \tilde{\phi}' + \lambda \tilde{\phi} = 0, \quad \tilde{\phi}'(0) = \tilde{\phi}'(\pi R) = 0. \quad (\text{B.2})$$

As this equation still contains first derivatives of delta functions, we take the regularized form of the delta function and its derivative to be

$$\delta_\rho(y) = \begin{cases} \frac{1}{\rho} \left(1 - \frac{|y|}{\rho} \right) & |y| < \rho, \\ 0 & |y| > \rho, \end{cases} \quad \delta'_\rho(y) = \begin{cases} -\frac{1}{\rho^2} \text{sgn}(y) & |y| < \rho, \\ 0 & |y| > \rho. \end{cases} \quad (\text{B.3})$$

Employing this regularization means that we have to solve three differential equations with the appropriate (continuously differentiable) matching and boundary conditions. By introducing somewhat more elaborate notation the three cases can be treated simultaneously.

$$\tilde{\phi}'' + 2\mu_{(a)} \tilde{\phi}' + \lambda \tilde{\phi} = 0, \quad \mu_{(a)} = g_5 q_b \left(\frac{1}{2} \xi_0 - a \frac{\xi_0''}{\rho^2} \right), \quad (\text{B.4})$$

where $a = +, 0, -$ refers to the regions $[0, \rho[$, $]\rho, \pi R - \rho[$ and $]\pi R - \rho, \pi R]$, respectively. The solutions of the equation are, of course, exponentials $\tilde{\phi}(y) = \exp(\alpha_{(a)}^b y)$ with the exponents

$$\alpha_{(a)}^b = -\mu_{(a)} + b \sqrt{\mu_{(a)}^2 - \lambda}, \quad b = \pm, \quad \alpha_{(a)}^+ \alpha_{(a)}^- = \lambda. \quad (\text{B.5})$$

For $\alpha_{(0)}^\pm$ we often use the short hand $\alpha^\pm$. Notice that if $\xi_0'' = 0$, for all a we have that $\alpha_{(a)}^b = \alpha^b$. The solution, with the boundary conditions (B.2) taken into account, can be represented as

$$\tilde{\phi}(y) = \begin{cases} A_{(+)} \left(\alpha_{(+)}^- e^{\alpha_{(+)}^+ y} - \alpha_{(+)}^+ e^{\alpha_{(+)}^- y} \right), & 0 < y < \rho, \\ A^+ e^{\alpha^+ y} + A^- e^{\alpha^- y}, & \rho < y < \pi R - \rho, \\ A_{(-)} \left(\alpha_{(-)}^- e^{\alpha_{(-)}^+ (y - \pi R)} - \alpha_{(-)}^+ e^{\alpha_{(-)}^- (y - \pi R)} \right), & \pi R - \rho < y < \pi R. \end{cases} \quad (\text{B.6})$$

By gluing the solutions together at the two boundaries between the three regions in a continuously differentiable fashion, allows one to express $A^\pm$ linearly in terms of either $A_{(+)}$ or $A_{(-)}$. Therefore, in

the ratio $-A^+/A^-$ these constants drop out and a consistency condition resulting from the boundary conditions is obtained. This consistency requirement

$$e^{(\alpha^+ - \alpha^-)(\pi R - 2\rho)} = \frac{1 - \alpha^- H_{(-)}}{1 - \alpha^+ H_{(-)}} \frac{1 - \alpha^+ H_{(+)}}{1 - \alpha^- H_{(+)}} , \quad H_{(a)} = \frac{1}{\alpha_{(a)}^+ \alpha_{(a)}^-} \frac{\alpha_{(a)}^- e^{a\alpha_{(a)}^+ \rho} - \alpha_{(a)}^+ e^{a\alpha_{(a)}^- \rho}}{e^{a\alpha_{(a)}^+ \rho} - e^{a\alpha_{(a)}^- \rho}} , \quad (\text{B.7})$$

leads to the quantization of the eigenvalue λ .

As a consistency check of this equation we observe that in the case $\xi_0'' = 0$, dividing the interval $[0, \pi R]$ into three regions is not necessary and therefore the result of the spectrum should be independent of ρ . Indeed, in this case we find that the quantization condition (B.7) reduces to

$$e^{(\alpha^+ - \alpha^-)\pi R} = 1 \quad \Rightarrow \quad \lambda_n = \frac{1}{4}(g_5 q_b \xi_0)^2 + \frac{n^2}{R^2}, \quad n \in \mathbb{N}, \quad (\text{B.8})$$

for any ρ . After the implication sign we used eqs. (B.4) and (B.5) and that eq. (B.8) has solutions only if $\lambda > \frac{1}{4}(g_5 q_b \xi_0)^2$. (For equality, the general solution is not of the form of a sum of two exponentials, and does not allow for non-trivial solutions of the boundary conditions. This excludes the would be zero mode $n = 0$, which we have already treated in detail in section 5.2.)

For $\xi_0'' \neq 0$, we are interested in the limit $\rho \rightarrow 0$ where the delta function and its derivative is reproduced. Although the asymptotics

$$H_{(\pm)} \rightarrow \begin{cases} \pm 0 & \xi_0'' q_b > 0, \\ \mp \infty & \xi_0'' q_b < 0, \end{cases} \quad (\text{B.9})$$

of the functions $H_{(\pm)}$ is very different for the cases $\text{sgn}(\xi_0'' q_b) = \pm$, in the limit of $\rho \rightarrow 0$ of (B.7) leads to the same quantization condition (B.8). (Here ± 0 denotes where the limit 0 is reached from above or below.) With this result we have shown, that it does not matter for the spectrum whether the term with ξ_0'' is present or not, provided that the branes are infinitely thin ($\rho \rightarrow 0$); as is dictated by the orbifolding.

For the odd bosonic states, we can perform a completely analogous analysis, therefore we just quote the main results. The boundary conditions for odd fields $\tilde{\phi}(0) = \tilde{\phi}(\pi R) = 0$, is implemented by the solution

$$\tilde{\phi}(y) = \begin{cases} A_{(+)} \left(e^{\alpha_{(+)}^+ y} - e^{\alpha_{(+)}^- y} \right), & 0 < y < \rho, \\ A^+ e^{\alpha^+ y} + A^- e^{\alpha^- y}, & \rho < y < \pi R - \rho, \\ A_{(-)} \left(e^{\alpha_{(-)}^+ (y - \pi R)} - e^{\alpha_{(-)}^- (y - \pi R)} \right), & \pi R - \rho < y < \pi R. \end{cases} \quad (\text{B.10})$$

This gives rise to the quantization equation of the form as in the even case, see (B.7), except the functions $H_{(\pm)}$ are different:

$$e^{(\alpha^+ - \alpha^-)(\pi R - 2\rho)} = \frac{1 - \alpha^- H_{(-)}}{1 - \alpha^+ H_{(-)}} \frac{1 - \alpha^+ H_{(+)}}{1 - \alpha^- H_{(+)}} , \quad H_{(a)} = \frac{e^{a\alpha_{(a)}^+ \rho} - e^{a\alpha_{(a)}^- \rho}}{\alpha_{(a)}^- e^{a\alpha_{(a)}^+ \rho} - \alpha_{(a)}^+ e^{a\alpha_{(a)}^- \rho}} . \quad (\text{B.11})$$

However the asymptotics is identical to (B.9), so that the same conclusion is reached: the spectrum is independent of ξ_0'' as long as $\rho \rightarrow 0$, and is given by (B.8).

C Characteristic classes and anomalies

It is well-known (see for example [28, 29]) that the form of possible counter terms for anomalies is dictated by characteristic classes. For chiral (non-Abelian) anomalies in $2n$ dimensions, the starting point is the formal $2(n+1)$ form

$$\Omega_{2(n+1)}(R, F) = \hat{A}(R) \text{ch}(F) \Big|_{2n+2}, \quad (C.1)$$

$$\hat{A}(R) = \text{tr} \frac{R/4\pi}{\sinh(R/4\pi)}, \quad \text{ch}(F) = \text{tr} \exp \frac{iF}{2\pi}.$$

Here $\text{ch}(F)$ denotes the Chern character of the field strength 2-form F of a gauge connection A_M ; and $\hat{A}(R)$ the roof genus of the curvature 2-form R . In this work we focus primarily on gauge anomalies, neglecting the gravitational and mixed anomalies, therefore we do not take the roof genus contribution into account from this point onwards (except when we comment on mixed gauge-gravitational-like anomaly contributions).

By the descent equations the following $2n+1$ form $\Omega_{2n+1}(A)$ and $2n$ form $\Omega_{2n}^1(A; \Lambda)$ are defined

$$d\Omega_{2n+1}(A) = \Omega_{2(n+1)}(F), \quad \delta_\Lambda \Omega_{2n+1}(A) = d\Omega_{2n}^1(A; \Lambda), \quad (C.2)$$

with Λ the infinitesimal parameter of a gauge transformation $\delta_\Lambda A = d\Lambda + [\Lambda, A]$.

The effective (Euclidean) action is obtained by integrating out the fermions ψ

$$e^{-\Gamma(A)} = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S}, \quad S = - \int dx \bar{\psi} \not{D}(A) \psi. \quad (C.3)$$

In the covariant derivative $D_M = \partial_M + A_M$ the gauge field A_M resides. There is no local gauge anomaly if for any local gauge parameter $\Lambda(x)$

$$\delta_\Lambda \Gamma(A) = \Gamma(A + \delta_\Lambda A) - \Gamma(A) = 0; \quad (C.4)$$

otherwise there is a gauge anomaly. For Minkowski-spaces gauge anomalies can only occur in even $2n$ dimensions. The general form of this anomaly is determined by the Wess-Zumino consistency conditions [25] for anomalies and the connection with the Atiyah-Singer index [30] to be

$$\delta_\Lambda \Gamma(A) = 2\pi i \int_{S^{2n}} \Omega_{2n}^1(A; \Lambda). \quad (C.5)$$

Here $2n$ dimensional Minkowski space has been replaced by its Euclidean analog and a point at infinity is added to obtain the topology of a $2n$ dimensional sphere S^{2n} .

In odd $(2n+1)$ dimensional Minkowski spaces no gauge anomalies can appear. The reason for this is simply that integrals over closed surfaces of closed forms are zero. The Wess-Zumino consistency conditions imply that the only possible form of the anomaly odd dimensions is

$$\delta_\Lambda \Gamma(A) = 2\pi i \int_{S^{2n+1}} d\Omega_{2n}^1(A; \Lambda). \quad (C.6)$$

But this vanishes identically, as a closed form is integrated over a closed surface.

D Gauge invariance of the parity anomaly counter term

In this appendix we show, using topological methods, that the parity anomaly counter term Γ_{PAC} is not necessarily gauge invariant [19], if the gauge group G contains a factor of $SU(n)$ (with $n \geq 3$) or $USp(2n)$, or at least one $U(1)$ factor and a simple compact Lie group.

Let us first assume that the group G is connected. Under a large gauge transformation the parity counter term variation equals the winding number of the gauge transformation in the group

$$\Gamma_{PAC}(^g A) = -\pi i \frac{2}{5!} \left(\frac{i}{2\pi} \right)^3 \int_{S^4 \times S^1} \text{tr}(g^{-1} dg)^5, \quad (\text{D.1})$$

which can be non-zero if the gauge transformation is a non-trivial map from $S^4 \times S^1$ or S^5 into the gauge group G .⁷ Therefore there are two possibilities: either both $\pi_4(G)$ and $\pi_1(G)$, or $\pi_5(G)$ have to be non-trivial. According to the table in [31]

$$\pi_1(G) = \begin{cases} \mathbb{Z} & U(1), \\ \mathbb{Z}_2 & SO(n \geq 3), \\ 0 & \text{others}, \end{cases} \quad \pi_4(G) = \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_2 & SO(4), Spin(4), \\ \mathbb{Z}_2 & SU(2), SO(3), Spin(5), SO(5), USp(2n), \\ 0 & \text{others} \end{cases} \quad (\text{D.2})$$

the only groups that have both these homotopy groups non-trivial are $SO(n)$, $n = 3, 4, 5$. However, the generators of $SO(n)$ are anti-symmetric hence the symmetric trace over the generators (D_{ijk}) is zero. The second possibility is a non-trivial π_5 homotopy group

$$\pi_5(G) = \begin{cases} \mathbb{Z} & SU(n \geq 3), \\ \mathbb{Z}_2 & USp(2n). \end{cases} \quad (\text{D.3})$$

We see that the parity anomaly counter term may be gauge non-invariant for all $USp(2n)$ groups and for $SU(n)$ groups with $n \geq 3$.

Next, we consider the case that G consists of several connected factors. So let $G = G_1 \times G_4$ where the group G_1 is a connected group and G_4 may contain various disconnected factors. By taking different connected factors for G_1 and repeating the arguments which are presented below, all possibilities can be found in this way. The field strength of the G_1 factor is denoted by F_1 and the field strength for the G_4 is denoted by F_4 . The relevant characteristic class Ω_6 then reads (see (C.1))

$$\Omega_6 = \left(\frac{i}{2\pi} \right)^3 \left[\frac{1}{6} \text{tr} F_1^3 + \frac{1}{2} \text{tr} F_1 \text{tr} F_4^2 + \frac{1}{2} \text{tr} F_1^2 \text{tr} F_4 + \frac{1}{6} \text{tr} F_4^3 \right]. \quad (\text{D.4})$$

We can disregard the first and last terms in this expression: the first term we have already treated, while for the last term we simply repeat the analysis we are describing with $G = G_4$. From the remaining two 6-forms we obtain four different 5-forms, but since they are integrated over a manifold without boundary, we only need those two of them ($\text{tr} A_1 \text{tr} F_4^2$ and $\text{tr} F_1^2 \text{tr} A_4$) which can be readily integrated over $S^1 \times S^4$. These terms can give non-vanishing contributions to the gauge transformation of the parity counter term

$$\Gamma_{PAC}(^g A) \supset -\pi i \frac{i}{2\pi} \int_{S^1} \text{tr}(g_1^{-1} dg_1) \frac{-1}{8\pi^2} \int_{S^4} \text{tr} F_4^2 - \pi i \frac{-1}{8\pi^2} \int_{S^4} \text{tr} F_1^2 \frac{i}{2\pi} \int_{S^1} \text{tr}(g_4^{-1} dg_4). \quad (\text{D.5})$$

⁷ S^5 appears when we consider gauge transformations satisfying the condition $g(x, 0) = g(x, 2\pi R) = g$ for some constant g . In such a case $S^1 \times S^4$ becomes topologically equivalent to S^5

We only have to focus on the first term, since the second term will be taken into account when we take G_1 to be one of the other factors in G . Thus, for gauge violation of the parity counter term, we need that both integrals of the first term are non-zero. The second integral gives the instanton number, which is classified by $\pi_3(G_4)$ which is $\mathbb{Z}$ for any simple connected Lie group. The first integral of the first term is the winding number of the gauge transformation that maps S^1 in the group G_1 , classified by the fundamental group $\pi_1(G_1)$ given in (D.2). But as the generators of the $SO(n \geq 3)$ groups are, traceless because of their anti-symmetry, only the $U(1)$ can give rise to non-vanishing contributions.

Since, we need gauge invariance for any background gauge field configuration, the instanton number is arbitrary. Gauge invariance of the parity counter term is also maintained if it changes by an integer multiple of $2\pi i$ as the effective action remains invariant.

Notice that if gravitational interactions are also taken into account, also the gravitational instanton can appear in (D.5) so that also the gauge groups $G = U(1)$ can produce a non-gauge invariant parity counter term. For this it is sufficient that

$$\int_{S^1} \frac{i}{2\pi} \text{tr}(g_1^{-1} dg_1) = 1 \bmod 2, \quad (\text{D.6})$$

for some large gauge transformations.

References

- [1] S. Groot Nibbelink, H. P. Nilles and M. Olechowski, arXiv:hep-th/0203055, to appear in Phys. Lett. B.
- [2] D. M. Ghilencea, S. Groot Nibbelink and H. P. Nilles, Nucl. Phys. B **619** (2001) 385 [arXiv:hep-th/0108184].
- [3] R. Barbieri, L. J. Hall and Y. Nomura, arXiv:hep-ph/0110102.
- [4] C. A. Scrucca, M. Serone, L. Silvestrini and F. Zwirner, Phys. Lett. B **525** (2002) 169 [arXiv:hep-th/0110073].
- [5] R. Jackiw and C. Rebbi, Phys. Rev. D **13** (1976) 3398
- [6] N. Arkani-Hamed, T. Gregoire and J. Wacker, JHEP **0203** (2002) 055 [arXiv:hep-th/0101233].
- [7] D. E. Kaplan and T. M. Tait, JHEP **0111** (2001) 051 [arXiv:hep-ph/0110126].
- [8] N. Arkani-Hamed, A. G. Cohen and H. Georgi, Phys. Lett. B **516** (2001) 395 [arXiv:hep-th/0103135].
- [9] L. Pilo and A. Riotto, arXiv:hep-th/0202144.
- [10] R. Barbieri, R. Contino, P. Creminelli, R. Rattazzi and C. A. Scrucca, arXiv:hep-th/0203039.
- [11] E. A. Mirabelli and M. E. Peskin, Phys. Rev. D **58** (1998) 065002 [hep-th/9712214].
- [12] P. Horava and E. Witten, Nucl. Phys. B **460** (1996) 506 [arXiv:hep-th/9510209].

- [13] M. Zucker, “Off-shell supergravity in five dimensions and supersymmetric brane world scenarios”, PhD thesis Bonn University, BONN-IR-2000-10, ISSN-017208741
M. Zucker, JHEP **0008** (2000) 016 [hep-th/9909144].
- [14] E. Bergshoeff, R. Kallosh and A. Van Proeyen, JHEP **0010** (2000) 033 [hep-th/0007044].
- [15] W. Fischler, H. P. Nilles, J. Polchinski, S. Raby and L. Susskind, Phys. Rev. Lett. **47** (1981) 757.
- [16] L. Alvarez-Gaume and P. Ginsparg, Annals Phys. **161** (1985) 423 [Erratum-ibid. **171** (1985) 233].
- [17] E. Witten, Commun. Math. Phys. **100** (1985) 197.
- [18] E. Witten, Phys. Lett. B **117** (1982) 324.
- [19] L. Alvarez-Gaume, S. Della Pietra and G. W. Moore, Annals Phys. **163** (1985) 288.
- [20] M. F. Atiyah, V. K. Patodi and I. M. Singer, Math. Proc. Cambridge Phil. Soc. **79** (1980) 71.
- [21] Y. Kawamura, Prog. Theor. Phys. **103** (2000) 613 [arXiv:hep-ph/9902423].
- [22] Y. Kawamura, Prog. Theor. Phys. **105** (2001) 999 [arXiv:hep-ph/0012125].
- [23] A. Hebecker and J. March-Russell, Nucl. Phys. B **625** (2002) 128 [arXiv:hep-ph/0107039].
- [24] H. D. Kim, J. E. Kim and H. M. Lee, arXiv:hep-th/0204132.
- [25] J. Wess and B. Zumino, Phys. Lett. B **37** (1971) 95.
- [26] N. Arkani-Hamed and M. Schmaltz, Phys. Rev. D **61** (2000) 033005 [arXiv:hep-ph/9903417].
- [27] H. Georgi, A. K. Grant and G. Hailu, Phys. Rev. D **63** (2001) 064027 [arXiv:hep-ph/0007350].
- [28] M. Nakahara, “Geometry, Topology And Physics,” *Bristol, UK: Hilger (1990) 505 p. (Graduate student series in physics)*.
- [29] L. Alvarez-Gaume, “An Introduction To Anomalies,” HUTP-85/A092 *Lectures given at Int. School on Mathematical Physics, Erice, Italy, Jul 1-14, 1985*.
- [30] M. F. Atiyah and I. M. Singer, Annals Math. **87** (1968) 484, Annals Math. **87** (1968) 546.
- [31] S. Weinberg, “The quantum theory of fields. Vol. 2: Modern applications,” *Cambridge, UK: Univ. Pr. (1996) 489 p, appendix B*.